

Tabel van Laplace getransformeerden

No.	$f(t) = L^{-1}[F(s)]$	$F(s) = L[f(t)] = \int_0^\infty e^{-st} f(t) dt$
(l.1)	1	$\frac{1}{s}, s > 0$
(l.2)	e^{at}	$\frac{1}{s-a}, s > a$
(l.3)	$\sin at$	$\frac{a}{s^2 + a^2}, s > 0$
(l.4)	$t^n, n \in \mathbb{N}$	$\frac{n!}{s^{n+1}}, s > 0$
(l.5)	$t^p, p > -1$	$\frac{\Gamma(p+1)}{s^{p+1}}, s > 0$
(l.6)	$\cos at$	$\frac{s}{s^2 + a^2}, s > 0$
(l.7)	$\sinh at$	$\frac{a}{s^2 - a^2}, s > a $
(l.8)	$\cosh at$	$\frac{s}{s^2 - a^2}, s > a $
(l.9)	$e^{at} \sin bt$	$\frac{b}{(s-a)^2 + b^2}, s > a$
(l.10)	$e^{at} \cos bt$	$\frac{s-a}{(s-a)^2 + b^2}, s > a$
(l.11)	$t^n e^{at}, n \in \mathbb{N}$	$\frac{n!}{(s-a)^{n+1}}, s > a$
(l.12)	$u_a(t)$	$\frac{e^{-as}}{s}, s > 0$
(l.13)	$u_a(t)f(t-a)$	$e^{-as} F(s)$
(l.14)	$e^{at} f(t)$	$F(s-a)$
(l.15)	$f(at)$	$\frac{1}{a} F\left(\frac{s}{a}\right), a > 0$
(l.16)	$\int_0^t f(t-\tau) g(\tau) d\tau$	$F(s) G(s)$
(l.17)	$\delta(t-a)$	e^{-as}
(l.18)	$f^{(n)}(t)$	$s^n F(s) - s^{n-1} f(0) - \dots - f^{(n-1)}(0)$
(l.19)	$(-t)^n f(t)$	$F^{(n)}(s)$

Formuleblad behorend bij Analyse door J.H.J. Almering e.a.

Enkele goniometrische formules

No.

- (g.1) $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta.$
- (g.2) $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta.$
- (g.3) $\sin \alpha + \sin \beta = 2 \sin \frac{1}{2}(\alpha + \beta) \cos \frac{1}{2}(\alpha - \beta).$
- (g.4) $\cos \alpha + \cos \beta = 2 \cos \frac{1}{2}(\alpha + \beta) \cos \frac{1}{2}(\alpha - \beta).$
- (g.5) $\cos \alpha - \cos \beta = -2 \sin \frac{1}{2}(\alpha + \beta) \sin \frac{1}{2}(\alpha - \beta).$
- (g.6) $\sin^2 \alpha = \frac{1}{2}(1 - \cos 2\alpha).$
- (g.7) $\cos^2 \alpha = \frac{1}{2}(1 + \cos 2\alpha).$
- (g.8) $1 + \tan^2 \alpha = \frac{1}{\cos^2 \alpha}.$
- (g.9) $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}.$

Standaard Taylorontwikkelingen

No.

- (s.1) $e^x = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} + O(x^{n+1}).$
 - (s.2) $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + O(x^{2n+3}).$
 - (s.3) $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^n \frac{x^{2n}}{(2n)!} + O(x^{2n+2}).$
 - (s.4) $\frac{1}{1-x} = 1 + x + x^2 + \dots + x^n + O(x^{n+1}).$
 - (s.5) $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^n \frac{x^{n+1}}{n+1} + O(x^{n+2}).$
 - (s.6) $\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots + (-1)^n \frac{x^{2n+1}}{2n+1} + O(x^{2n+3}).$
 - (s.7) $(1+x)^a = 1 + ax + \frac{a(a-1)}{2!} x^2 + \dots + \frac{a}{n} x^n + O(x^{n+1}), a \in \mathbb{R}.$
- Hierbij is $\frac{a}{n} = \frac{a(a-1)\dots(a-n+1)}{n!}$ als $n > 0$ en $\frac{a}{0} = 1.$

Integraaltabel

No.

$$(t.1) \quad \int x^a dx = \frac{1}{a+1} x^{a+1} \quad (a \in \mathbb{R} \setminus \{-1\}).$$

$$(t.2) \quad \int \frac{dx}{x} = \ln|x|.$$

$$(t.3) \quad \int \frac{f'(x)}{f(x)} dx = \ln|f(x)|.$$

$$(t.4) \quad \int e^x dx = e^x.$$

$$(t.5) \quad \int a^x dx = \frac{1}{\ln a} a^x \quad (a \in \mathbb{R}^+ \setminus \{1\}).$$

$$(t.6) \quad \int \sin x dx = -\cos x.$$

$$(t.7) \quad \int \cos x dx = \sin x.$$

$$(t.8) \quad \int \frac{dx}{\sin^2 x} = -\cot x.$$

$$(t.9) \quad \int \frac{dx}{\cos^2 x} = \tan x.$$

$$(t.10) \quad \int \tan x dx = -\ln|\cos x|.$$

$$(t.11) \quad \int \frac{dx}{\sin x} = \ln|\tan \frac{x}{2}| = \ln \frac{1 - \cos x}{|\sin x|} = \ln \frac{|\sin x|}{1 + \cos x}.$$

$$(t.12) \quad \int \frac{dx}{\cos x} = \ln|\tan(\frac{x}{2} + \frac{\pi}{4})| = \ln \frac{1 + \sin x}{|\cos x|} = \ln \frac{|\cos x|}{1 - \sin x}.$$

$$(t.13) \quad \int \sin^p x \cos^q x dx \quad (p, q \in \mathbb{N})$$

p oneven, stel $\cos x = t$.
 q oneven, stel $\sin x = t$.
 p en q even, voer dubbele hoek in met behulp
 van $\cos^2 x = \frac{1 + \cos 2x}{2}$ of $\sin^2 x = \frac{1 - \cos 2x}{2}$.

$$(t.14) \quad \int \sin mx \sin nx dx = \frac{\sin(m-n)x}{2(m-n)} - \frac{\sin(m+n)x}{2(m+n)} \quad (m^2 \neq n^2).$$

$$(t.15) \quad \int \cos mx \cos nx dx = \frac{\sin(m-n)x}{2(m-n)} + \frac{\sin(m+n)x}{2(m+n)} \quad (m^2 \neq n^2).$$

$$(t.16) \quad \int \sin mx \cos nx dx = -\frac{\cos(m-n)x}{2(m-n)} - \frac{\cos(m+n)x}{2(m+n)} \quad (m^2 \neq n^2).$$

$$(t.17) \quad \int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx) \quad (a^2 + b^2 \neq 0).$$

$$(t.18) \quad \int e^{ax} \cos bx dx$$

$$= \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx) \quad (a^2 + b^2 \neq 0).$$

$$(t.19) \quad \int \frac{dx}{a^2 + b^2 x^2}$$

$$= \frac{1}{ab} \arctan \frac{bx}{a} \quad (a, b \in \mathbb{R}^+).$$

$$(t.20) \quad \int \frac{dx}{a^2 - b^2 x^2}$$

$$= \frac{1}{2ab} \ln \left| \frac{a+bx}{a-bx} \right| \quad (a, b \in \mathbb{R}^+).$$

$$(t.21) \quad \int \frac{dx}{\sqrt{a^2 - x^2}}$$

$$= \arcsin \frac{x}{a} \quad (a \in \mathbb{R}^+).$$

$$(t.22) \quad \int \frac{dx}{\sqrt{a^2 + x^2}}$$

$$= \ln(x + \sqrt{x^2 + a^2}) \quad (a \in \mathbb{R}^+).$$

$$(t.23) \quad \int \frac{dx}{\sqrt{x^2 - a^2}}$$

$$= \ln|x + \sqrt{x^2 - a^2}| \quad (a \in \mathbb{R}^+).$$

$$(t.24) \quad \int \sqrt{a^2 - x^2} dx$$

$$= \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin \frac{x}{a} \quad (a \in \mathbb{R}^+).$$

$$(t.25) \quad \int \sqrt{a^2 + x^2} dx$$

$$= \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \ln(x + \sqrt{a^2 + x^2}) \quad (a \in \mathbb{R}^+).$$

$$(t.26) \quad \int \sinh x dx$$

$$= \cosh x.$$

$$(t.27) \quad \int \cosh x dx$$

$$= \sinh x.$$

$$(t.28) \quad \int_0^a \sqrt{a^2 - x^2} dx$$

$$= \frac{\pi a^2}{4} \quad (a \in \mathbb{R}^+).$$

$$(t.29) \quad \int_0^{\frac{\pi}{2}} \sin^p x dx = \int_0^{\frac{\pi}{2}} \cos^p x dx$$

$$= \frac{p-1}{p} \frac{p-3}{p-2} \dots \frac{4}{5} \frac{2}{3} \text{ voor } p = \text{oneven } \geq 3,$$

$$= \frac{p-1}{p} \frac{p-3}{p-2} \dots \frac{3}{4} \frac{1}{2} \text{ voor } p = \text{even } \geq 2.$$

$$(t.30) \quad \int_0^{\infty} e^{-x} x^n dx$$

$$= n! \quad (n \in \mathbb{N}).$$

$$(t.31) \quad \int_0^1 x^n (1-x)^m dx$$

$$= \frac{(n!)(m!)}{(m+n+1)!} \quad (n, m \in \mathbb{N}).$$